

CHEST TRAUMA

Delayed respiratory deterioration post blunt chest trauma Immersive scenario

Facilitator resource kit





Queensland Trauma Education

The resources developed for Queensland Trauma Education are designed for use in any Queensland Health facility that cares for patients who have been injured as a result of trauma. Each resource can be modified by the facilitator and scaled to the learners needs as well as the environment in which the education is being delivered, from tertiary to rural and remote facilities.

Developed by

Alison Blunt, Advanced Physiotherapist – Princess Alexandra Hospital

Reviewed by

Angelka Opie, Nurse Educator – CSDS, MNHHS

Peter Thomas, Consultant Physiotherapist – Royal Brisbane and Women's Hospital Fiona Jennings, Nurse Practitioner Trauma Service – Princess Alexandra Hospital Dr Frances Williamson, Staff Specialist Emergency Physician – Metro North Health Kimberly Ballinger, Simulation Educator – Clinical Skills Development Service Education Working Group, Statewide Trauma Clinical Network – Clinical Excellence Queensland

Queensland Trauma Education

Chest Trauma – Delayed respiratory deterioration post blunt chest trauma: Immersive scenario – Facilitator resource kit
Version 2.0

Published by the State of Queensland (Clinical Skills Development Service), 2024



This document is licensed under a Creative Commons Attribution 3.0 Australia licence. To view a copy of this licence, visit https://creativecommons.org/licenses/by/3.0/au.

© State of Queensland (Metro North Hospital and Health Service through the Clinical Skills Development Service) 2021

You are free to copy, communicate, and adapt the work, as long as you attribute the Metro North Hospital and Health Service through the Clinical Skills Development Service. For more information, please contact Clinical Skills Development Service, Royal Brisbane and Women's Hospital, Herston, Queensland +61 3646 6500, CSDS-Admin@csds.qld.edu.au.

An electronic version of this document is available via csds.qld.edu.au/qte

Disclaimer: The content presented in this publication is distributed by the Queensland Government as an information source only. The State of Queensland makes no statements, representations or warranties about the accuracy, completeness or reliability of any information contained in this publication. The State of Queensland disclaims all responsibility and all liability (including without limitation for liability in negligence) for all expenses, losses, damages and costs you might incur as a result of the information being inaccurate or incomplete in any way, and for any reason reliance was placed on such information.

About this training resource kit

This resource kit provides healthcare workers with useful resources and key practice points to assist them to identify and manage respiratory complications in patients following blunt chest trauma injuries.

National Safety and Quality Health Service (NSQHS) Standards













Target audience

Ward-based nursing and physiotherapy clinicians.

Duration

45-60 minutes (set-up, scenario and debriefing).

Group size

4-6 participants (or team composition applicable to local area).

Learning objectives

By the end of this session the participant will be able to:

- Identify patient and injury-related factors that place a patient at greater risk of delayed respiratory complications post chest trauma.
- Identify three key signs of impending respiratory deterioration poorly controlled pain, impaired inspiration, and impaired cough.
- Demonstrate the skill of teaching a patient the correct incentive spirometry technique.

Facilitation guide

- 1. OPTIONAL: Utilise associated PowerPoint presentation to provide participants with clinical management of blunt chest trauma prior to delivery of immersive scenario.
- 2. Facilitator to discuss the pre-simulation briefing and deliver the immersive scenario on blunt chest trauma.
- 3. Utilise the supporting documents to maximise the learning throughout immersive scenario
- 4. Utilise the debriefing guide to evaluate participant performance and provide feedback.

Supporting documents

The following supporting documents are provided for this immersive scenario:

- 1. Structured assessment and specific management
- 2. Pre-simulation briefing poster
- CXR1: Day 1
 CXR2: Day 2
- 5. Flowchart: Blunt Chest Trauma
- 6. PIC Score

Simulation event

This section contains the following:

- 1. Immersive scenario
- 2. Resource requirements
- 3. Handover card
- 4. Scenario progression
 - a. State 1: Initial assessment / recognition of deterioration
 - b. State 2: Ongoing management / secondary assessment
 - c. State 3: Reassessment
- 5. Debriefing guide

Immersive scenario

Туре	Immersive scenario	
Target audience	Ward-based nursing and physiotherapy staff.	
Overview	69 year old patient who is day 2 post a 2m fall from a ladder at home. Injuries sustained include fractured ribs R) 3-7 with no flail, right fractures to mid shaft tibia and fibula, and minor head abrasions.	
	CT head, chest and pelvis were NAD except for the above rib fractures, a R) pneumothorax, and bilateral pulmonary contusions.	
	A R) intercostal catheter (ICC) was inserted in the emergency department on admission to their local hospital.	
	Yesterday they underwent ORIF for the lower limb fractures and is now on the orthopaedic surgical ward.	
Learning objectives	 Identify patient and injury-related factors that place a patient at greater risk of delayed respiratory complications post chest trauma. 	
	 Identify three key signs of impending respiratory deterioration – poorly controlled pain, impaired inspiration, and impaired cough. 	
	 Demonstrate the skill of teaching a patient the correct incentive spirometry technique (optional). 	
Duration	45-60 minutes, including debrief.	

Resource requirements

Physical resources

Room setup	Hospital bed in ward environment	
Simulator/s	Simulated patient x1 (male or female)	
Simulator set up	 Hospital gown and TED stockings, lying supine in bed Moulage: Bruising to right chest, apical and basal chest drains attached to UWSD. Right leg in POP backslab or Moon boot. Dressing on forehead. 	
Clinical equipment	 NIBP, ECG and Sp02 monitoring. Nasal prongs attached to oxygen source. ICC and UWSD set up. Patient controlled analgesia (PCA), labelled "Morphine". PPE – hand sanitizer gloves. Incentive spirometer (if available). Fisher & Paykel – Airvo. High flow humidification device (Airvo), HFNP interface, circuit, water for irrigation. 	
Access	Left cubital 1 x IVC attached to PCA system	
Other	 Vital signs observations chart – QADDS or iEMR Patient Deterioration. PowerPoint slides. 	

Human resources

Faculty	2 facilitators (physiotherapist or nurse with debriefing experience) to take on roles of: • scenario commander/primary debrief • confederate nurse.	
Simulation coordinators	Standardised patient – facilitators to control simulated monitor.	
Confederates	Confederate nurse in room to give handover and responses.	
Other	Simulation technical or faculty staff to run the vital signs monitor (if available).	

Handover card

Handover from bedside nurse

This is Lesley, a 69 year old new admission to the orthopaedic ward overnight.

They were BIBA from home to the emergency department yesterday morning after an unwitnessed fall of approximately 2 metres from a ladder while attempting to trim trees. QAS reported the patient was lying on their right side with pain in the right chest, lower leg, and minor forehead abrasions but no loss of consciousness at the scene.

CT head, chest and pelvis were NAD except for fractured ribs R) 3-7 with no flail, a R) pneumothorax, and bilateral lung contusions. A R) intercostal catheter (ICC) was inserted in the emergency department on admission to their local hospital. Yesterday they underwent ORIF for the lower limb fractures. Post-operative weight bearing orders are for NWB on the right leg for six weeks.

Lesley has no significant PMHx, and is normally fit and well.

Lives with partner and two university-aged children.

Smoking history of 1 pack of cigarettes per day since age 22.

Lesley has slept poorly overnight due to high levels of pain and they needs to be frequently reminded to use the PCA. I am concerned because in the last two hours their respiratory rate has increased and SpO2 has decreased.

Scenario progression

	STATE 1: INITIAL ASSESSMENT / RECOGNITION OF DETERIORATION				
Vital sign	ıs	Script	Details	Expected actions	
ECG HR	Sinus 99 92% on	"My ribs hurt!!" "My pain is 6/10 at rest, and 10/10 (chest) when I try to cough." "I can't cough, I have phlegm stuck." "I haven't slept a wink in 2 days." Nurse "They keep forgetting to use their PCA." "I've had to turn the oxygen up from 2 to 3 litres/min in the last hour." "I have just given some paracetamol but it doesn't	Primary survey results A: Patent B: Breathing pattern looks laboured, using accessory	 Commence Primary Survey Assesses "Pain, Inspiration, Cough Evaluate pain levels – rest, movement. Evaluate inspiratory capacity – observe 	
SpO ₂	3LO ₂ via nasal prongs		muscles. Reduced chest wall movement right side. Unable to take full	 chest wall movement, auscultate, perform incentive spirometry (if available). Evaluate cough effectiveness – pain 	
BP/ART RR	159/99 25		inspiration or cough, pain inhibited. C: Well perfused	inhibited. Recognises "red flags" for potential deterioration.	
Temp	38.1		D: GCS E3V5M6 E: Lying slumped in bed with right leg elevated, in back slab or Moon boot. R) ICC minimal output, nil bubble or swing	 >3 fractured ribs age >55 years uncontrolled pain increased WOB, RR decreased SpO2 smoking history pulmonary contusion decreased mobility (leg ORIF). 	
		seem to be working yet." "They haven't yet mobilised out of bed because of the leg fractures" "The chest drain output was about 100mls in total overnight"		 Request Acute Pain Management Service/ medical review/notify team leader. Nurse administered bolus morphine. Refer to physiotherapy if not already involved. 	

	STATE 2: ONGOING MANAGEMENT / SECONDARY ASSESSMENT			
Vital sign	S	Script	Details	Expected actions
ECG	Sinus	Lesley	Secondary survey results	Secondary survey
HR	92	"My pain is a bit better, 5/10." "It's still a bit hard to breathe."	Moist sounding cough. ICC site visible, dressing and	Assess other injuries, order investigations.
SpO ₂	94% on 4LO ₂ nasal prongs	"It hurts when I cough." "I'm knackered."	connections intact. Left calf examination NAD. Pulses intact. Right calf immobilized in boot or backslab, but neurovascularly intact.	Investigations ☐ Septic screen – FBC, sputum m/c/s, CXR.
BP/ART	144/81	"I'm a bit concerned about this patient. I looked after them on the late shift yesterday and they seem a bit worse today."	Julian, var nou o vaccular, j macu	Management
RR	22		em on the late shift control c	Reposition high sitting in bed.Hourly deep breathing exercises
Temp	37.8		pending.	+/- incentive spirometry.
GCS	15			Commenced on high flow nasal oxygen therapy (e.g. flow rate 40L/min, Fi02 0.32 or similar).
				 Acute Pain Management Service review – consideration for regional block.
				☐ Medical/ICU review.

	STATE 3: REASSESSMENT			
Vital sign	S	Script	Details	Expected actions
ECG HR	Sinus 88	Person "That's a bit better	Assessment results Reported pain levels improved to 2/10	Assessment Pain, inspiratory capacity and
SpO ₂	96% on HFNP FiO ₂ 0.32	(breathing)." Nurse These are the CXR results (hand hard copies of Day 1 and Day 2 CXRs to participants to review)	rest, 4/10 with movement. Coughing more effectively. Incentive spirometry performance improved. WCC 15.1. CXR – as per image.	cough. Investigations □ Review CXR. Management □ Continue regular analgesia and deep breathing exercises.
BP/ART RR	134/79 20			
Temp	37.5 15			

Debriefing guide

Scenario objectives

- Identify factors that place a patient at greater risk of delayed respiratory complications post chest trauma.
- Demonstrate the bedside assessment of a patient with chest trauma, including estimation of pain, inspiratory capacity and cough effectiveness.
- Identify and act on signs of respiratory deterioration.
- Demonstrate the skill of teaching a patient the correct incentive spirometry technique (optional).

Example questions

Exploring diagnosis

- What chest trauma "red flags" for respiratory deterioration, if any, were present in this case scenario? Refer to flowcharts from Clinical Practice Guideline: Blunt chest trauma (supporting documents).
- What could be some of the pathologies responsible for Lesley's deterioration on day 2?
- What abnormalities did you see on the CXR?

Discussing management and teamwork

- What members of the multidisciplinary team are involved in the detection and management of respiratory deterioration?
- Why was an ICU review recommended for Lesley?
- What are some other possible options (other than a PCA) for improved pain management for Lesley?
- How might Lesley's other injuries have affected her presentation and management? (possible head injury - although serious pathology excluded on CT, the timing of her ORIF under general anaesthetic yesterday for her leg fractures).

Key moments

- Recognising the "red flags" for potential deterioration in Lesley's history and presentation.
- Alerting team leader/ medical team / requesting Acute Pain Management Service review.
- Implementing steps to improve Lesley's pain control and inspiratory capacity.

Acronyms and abbreviations

Term	Definition	
CXR	Chest x-ray	
GCS	Glasgow coma scale	
HFNP	High flow nasal prongs	
ICC	Intercoastal catheter	
ORIF	Open reduction and internal fixation	
QAS	Queensland Ambulance Service	
СТ	Computed tomography	
TED	Thrombo-embolus deterrent	
PCA	Patient controlled analgesia	
iEMR	Integrated electronic medical record	
POP	Plaster of paris	
UWSD	Underwater seal drain	
PPE	Personal protective equipment	
QADDS	Queensland adult deterioration detection system	
BIBA	Brought in by ambulance	
NAD	Nil abnormalities detected	
NWB	Non weight bearing	
PMHx	Past medical history	
FBC	Full blood count	
ICU	Intensive care unit	
FiO2	Fraction of inspired oxygen	
WCC	White cell count	

References

- Battle, C.E., Hutchings, H., James, K.. & Evans, P.A. (2013). The risk factors for the development of complications during the recovery phase following blunt chest wall trauma: a retrospective study. *Injury*, 44(9), 1171-6. https://doi.org/10.1016/j.injury.2012.05.019
- 2. Ludwig, C. & Koryllos, A. (2017). Management of chest trauma. *Journal of thoracic disease*, *9*(Suppl 3), S172–S177. https://doi.org/10.21037/jtd.2017.03.52
- 3. Sadler, C., Burgess, J., Dougherty, K., & Collins, J. (2019). Bedside incentive spirometry predicts risk of pulmonary complications in patients with rib fractures. *The American Surgeon*, *85*, 1051-1055. https://pubmed.ncbi.nlm.nih.gov/31638523/
- 4. Witt, C.E., & Bulger, E.M. (2017). Comprehensive approach to the management of the patient with multiple rib fractures: A review and introduction of a bundled rib fracture management protocol. *Trauma Surgery and Acute Care Open, 2*, 1-7. http://dx.doi.org/10.1136/tsaco-2016-000064

Share your feedback

Please complete our survey to help make Queensland Trauma Education better

The survey should take no more than 5 minutes to complete.

Scan the QR code or visit:

<u>Evaluation Form - Clinical Skills Development Service</u>
(csds.qld.edu.au)





Queensland Trauma Education

Chest Trauma – Delayed respiratory deterioration post blunt chest trauma: Immersive scenario

– Facilitator resource kit

Published by the State of Queensland (Clinical Skills Development Service), 2024

Visit csds.qld.edu.au/qte
Email CSDS-Admin@health.qld.gov.au
Phone +61736466500

