



Queensland
Trauma Education

PELVIC TRAUMA

Haemodynamically unstable pelvic trauma

Immersive scenario

Facilitator resource kit



CSDS



Clinical Skills Development Service



Queensland Trauma Education

The resources developed for Queensland Trauma Education are designed for use in any Queensland Health facility that cares for patients who have been injured as a result of trauma. Each resource can be modified by the facilitator and scaled to the learners needs as well as the environment in which the education is being delivered, from tertiary to rural and remote facilities.

Developed by

Dr Frances Williamson, Staff Specialist Emergency Physician – Metro North Health
Kimberly Ballinger, Simulation Educator – Clinical Skills Development Service

Reviewed by

Angelka Opie, Nurse Educator – CSDS, MNHHS

Education Working Group, Statewide Trauma Clinical Network – Clinical Excellence
Queensland

Queensland Trauma Education

Pelvic trauma – Haemodynamically unstable pelvic trauma: Immersive scenario – Facilitator resource kit

Version 2.0

Published by the State of Queensland (Clinical Skills Development Service), 2024



This document is licensed under a Creative Commons Attribution 3.0 Australia licence. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/3.0/au>.

© State of Queensland (Metro North Hospital and Health Service through the Clinical Skills Development Service) 2021

You are free to copy, communicate, and adapt the work, as long as you attribute the Metro North Hospital and Health Service through the Clinical Skills Development Service. For more information, please contact Clinical Skills Development Service, Royal Brisbane and Women's Hospital, Herston, Queensland +61 3646 6500, CSDS-Admin@csds.qld.edu.au.

An electronic version of this document is available via csds.qld.edu.au/qte

Disclaimer: The content presented in this publication is distributed by the Queensland Government as an information source only. The State of Queensland makes no statements, representations or warranties about the accuracy, completeness or reliability of any information contained in this publication. The State of Queensland disclaims all responsibility and all liability (including without limitation for liability in negligence) for all expenses, losses, damages and costs you might incur as a result of the information being inaccurate or incomplete in any way, and for any reason reliance was placed on such information.

About this training resource kit

This resource kit provides healthcare workers with the knowledge and skills to manage a patient with an open book pelvic injury who is haemodynamically unstable following a traumatic incident.

National Safety and Quality Health Service (NSQHS) Standards



Target audience

Emergency medical and nursing clinicians.

Duration

45-60 minutes.

Group size

4-6 participants (or team composition applicable to local area).

Learning objectives

By the end of this session the participant will be able to:

- perform a focussed clinical examination to assess a patient with a major pelvic injury
- identify types of pelvic injury that are associated with vascular injury and bleeding
- perform bedside interventions to aid haemorrhage management
- demonstrate early targeted management (may include retrieval services).

Facilitation guide

1. Facilitator to discuss the pre-simulation briefing and deliver the immersive simulation on haemodynamically unstable pelvic trauma.
2. Utilise the supporting documents to maximise learning throughout the immersive scenario.
3. Utilise the debrief guide to evaluate and support participant performance and provide feedback.

Supporting resources (in Printable resources)

1. Pre-simulation briefing poster
2. Structured assessment in trauma.
3. Venous blood gas

4. Pelvic Xray
5. Chest Xray
6. EFAST: RUQ/Morrisons: negative
7. EFAST: LUQ/splenorenal: negative
8. EFAST: Pelvis: negative
9. EFAST: Subxiphoid: negative

Specific management

- Recognition of open book pelvic injury.
- Application of pelvic binder.
- Haemostatic resuscitation.
- Identification of potential arterial bleeding and management options.

Simulation event

This section contains the following:

1. Immersive scenario
2. Resource requirements
3. Handover card
4. Scenario progression
 - a. State 1: Initial assessment
 - b. State 2: Ongoing management / secondary assessment
 - c. State 3: Management
5. Debriefing guide

Immersive scenario

| | |
|----------------------------|---|
| Type | Immersive scenario |
| Target audience | Emergency department medical and nursing clinicians |
| Overview | <p>A 30-year-old male patient is transported to the ED following an MBC 50 minutes ago. He is haemodynamically unstable, confused and pale. He is complaining of pain over R lower quadrant of his abdomen and hip.</p> <p>Participants are required to demonstrate the rapid recognition of haemodynamic instability and perform assessment focused on identification of source of bleeding and prioritise management.</p> |
| Learning objectives | <p>By the end of this session the participant will be able to:</p> <ul style="list-style-type: none">• perform the assessment of a haemodynamically unstable trauma patient to identify a major pelvic injury• apply external pelvic compressive device to aid haemorrhage management• implement haemostatic resuscitation strategy• demonstrate early targeted management. |
| Duration | 45-60 minutes, including debrief. |

Resource requirements

Physical resources

| | |
|---------------------------|---|
| Room setup | Resus bay in emergency |
| Simulator/s | 1 manikin - SimMan 3G or ALS simulator |
| Simulator set up | <ul style="list-style-type: none"> • Street clothes, lying supine • Cervical collar in situ • No pelvic binder • Moulage: anterior bruising across lower abdomen/pelvis |
| Clinical equipment | <ul style="list-style-type: none"> • Standard precautions PPE. • Standard resus bay equipment: monitors, resus trolley, infusion pumps, blood warmers. • Fluids/blood products: N/saline, Hartmann's, packed red blood cells/blood components • Medications: IV analgesia • Pelvic binder (available for application if requested by participants) |
| Access | 2 x IVC setups. 18g cannula R) ACF with empty N/S 0.9% 250ml bag, No IV sticker attached to L) arm |
| Other | ED chart and relevant paperwork |

Human resources

| | |
|--------------------------------|---|
| Faculty | 2 facilitators (doctor/nurse with debriefing experience) to take on roles of scenario commander and primary debriefer |
| Simulation coordinators | 1 for manikin set up and control |
| Confederates | 1 confederate in room, optional 1 confederate to provide QAS handover / radiographer / other team member |
| Other | Trauma team composition – 3-4 nurses and 3 doctors in room (or team composition applicable to local area) |

Handover card

Handover from ambulance officer

This is Sam Johnson. He is an otherwise healthy 30-year-old male with no known allergies. He was witnessed to be the rider of a motorbike vs truck approximately 50 minutes ago. He was thrown over the bike, impacting with the road about 5 meters away.

He was assisted by bystanders on scene until the first crew arrived but was reportedly never knocked out, was wearing a helmet and remains GCS 14 throughout care, being confused to place and time. He has no other focal neurological deficits, pupils 3mm and reactive bilaterally.

His other vital signs are: HR 120, BP 90/60mmHg, saturations 97% 6L HM and respiratory rate 28. He is complaining of pain in his R lower quadrant of his abdomen and R hip.

He has an 18g cannula in his R) ACF and we have given him 10mg IV morphine and 8mg IV ondansetron with minimal effect. He has also had 750mls NSaline IV and a cervical collar applied.

Many thanks for your ongoing care of Sam.

Scenario progression

| STATE 1: INITIAL ASSESSMENT | | | | |
|-----------------------------|-----------|---|---|--|
| Vital signs | | Script | Details | Expected actions |
| ECG | ST | Sam Moaning loudly, "I am in so much pain. Please help me." | Primary survey results A: Cervical collar, airway patent, nil anterior neck injury B: nil chest wall tenderness, nil crepitus, nil subcutaneous emphysema, equal breath sounds bilaterally C: nil external bleeding, poor perfusion peripherally, pelvis and lower abdo tenderness, bruising across lower abdomen/pelvis, scrotal/penile bruising D: GCS 14 (confused), nil neurological deficits E: temp 35.6 | Commence primary survey <ul style="list-style-type: none"> <input type="checkbox"/> Assess airway/breathing <ul style="list-style-type: none"> ○ optimise oxygenation and ventilation. <input type="checkbox"/> Assess circulation <ul style="list-style-type: none"> ○ recognise abnormality in circulation ○ gain further IV access. <input type="checkbox"/> Assess disability and expose patient. <input type="checkbox"/> Action call for help early. Identify resources available to local area. |
| HR | 120 | | | |
| SpO₂ | 97% 6L HM | | | |
| BP/ART | 90/60mmHg | | | |
| RR | 28 | | | |
| Temp | 35.6 | | | |
| BGL | 7 | | | |
| GCS | 14 E4V4M6 | | | |
| | | | | |

| STATE 2: ONGOING MANAGEMENT / SECONDARY ASSESSMENT | | | | |
|--|-----------|--|--|---|
| Vital signs | | Script | Details | Expected actions |
| ECG | ST | Sam Moaning, "Ahhhhh, my hip! What's happening?" | Secondary survey results If not examined above: Abdomen/pelvis: pelvis and lower abdo tenderness, bruising across lower abdomen/pelvis, scrotal/penile bruising No wounds to suggest compound injury Long bones: NAD Log roll: nil midline bony tenderness/bruising Results CXR: NAD Pelvic Xray: open book pelvic fracture FAST: negative | Secondary survey <ul style="list-style-type: none"> <input type="checkbox"/> Perform head to toe assessment <input type="checkbox"/> Identify major pelvic injury and circulation compromise <input type="checkbox"/> Arrange analgesia <input type="checkbox"/> Ensure oxygenation is adequate Initiate investigations <ul style="list-style-type: none"> <input type="checkbox"/> Bloods - trauma panel - FBE, chem20, group and hold, lipase, coags/ROTEM (if applicable) <input type="checkbox"/> Point of care tests: Hemocue, Istat CG4 (if applicable) <input type="checkbox"/> Bedside tests: UA, ECG, VBG <input type="checkbox"/> Imaging: CXR/Pelvic Xray & FAST Management <ul style="list-style-type: none"> <input type="checkbox"/> Apply pelvic binder and strap feet in internal rotation <input type="checkbox"/> Commence fluid resuscitation - blood/blood products as preference <input type="checkbox"/> Warm patient |
| HR | 120 | | | |
| SpO₂ | 97% 6L HM | | | |
| BP/ART | 80/50mmHg | | | |
| RR | 28 | | | |
| Temp | 35.6 | | | |
| BGL | 7 | | | |
| GCS | 14 | | | |

| STATE 3: MANAGEMENT | | | | |
|---------------------|------------|---|--|---|
| Vital signs | | Script | Details | Expected actions |
| ECG | ST | Sam "I still have pain but it feels a little better since the thing on my hips was put on..." | If pelvic binder applied and fluid resuscitation commenced - patient haemodynamics improve. If no pelvic binder or fluid resuscitation initiated - haemodynamic parameters worsen - Confederate can prompt team to perform both actions. | Assessment <input type="checkbox"/> Repeat primary survey Management <input type="checkbox"/> Consideration for CT abdo with improvement in haemodynamic status for IR (or OT if remains unstable) OR. <input type="checkbox"/> Liaise with tertiary referral facility for consultation. Referral to surgical/ortho team for ongoing operative management OR Consult RSQ for retrieval. |
| HR | 110 | | | |
| SpO ₂ | 97% 6l HM | | | |
| BP/ART | 100/60mmHg | | | |
| RR | 24 | | | |
| Temp | 35.6 | | | |
| BGL | 7 | | | |
| GCS | 14 | | | |
| | | | | |

Debriefing guide

Scenario objectives

- perform the assessment of a haemodynamically unstable trauma patient to identify a major pelvic injury
- apply external pelvic compressive device to aid haemorrhage management
- implement haemostatic resuscitation strategy demonstrate early targeted management.

Example questions

Exploring diagnosis

- Explain your thought process for the rapid assessment of the haemodynamically unstable trauma patient for identification of life-threatening injuries.
- What clinical findings aided in the identification of bleeding source?
- Do the radiological investigations and EFAST help you identify the type of bleeding - arterial or venous?
- What clinical features aided the classification of shock state for this patient into mild/moderate/severe?
- What are the signs of associated urethral injury with an open book pelvic fracture?

Discussing management

- What was your priority to manage the haemodynamic instability?
- What is a system for classification of pelvic fractures and how does this affect your management?
- Is interventional radiology available at your hospital? What processes need to occur to activate this service?
- How do you activate a massive transfusion/VHA guided resuscitation protocol?
- Are there challenges in placing an indwelling catheter in this patient?

Discussing teamwork / crisis resource management

- Calling for help early - did you have enough team members to simultaneously manage the patient?
- How do you prioritise the management to improve his haemodynamic state?

Key moments

- Rapid recognition of haemodynamic instability and assessment focused on identification of source of bleeding.
- Early application of pelvic binder with internal rotation of feet to aid haemorrhage control.
- Institution of haemostatic resuscitation.
- Decision making for disposition - CTA and IR vs OT.

Acronyms and abbreviations

| Term | Definition |
|-------|---|
| CTA | Computed tomography arterial |
| IR | Interventional radiology |
| OT | Operating theatre |
| VHA | Viscoelastic haemostatic assays |
| EFAST | Extended focused assessment with sonography in trauma |
| UA | Urinalysis |
| ECG | Electrocardiogram |
| NAD | No abnormality detected |
| RSQ | Retrieval Services Queensland |

References

1. Coccolini, F., Stahel, P. F., Montori, G., Biffl, W., Horer, T. M., Catena, F., Kluger, Y., Moore, E. E., Peitzman, A. B., Ivatury, R., Coimbra, R., Fraga, G. P., Pereira, B., Rizoli, S., Kirkpatrick, A., Leppaniemi, A., Manfredi, R., Magnone, S., Chiara, O., Solaini, L., ... Ansaloni, L. (2017). Pelvic trauma: WSES classification and guidelines. *World journal of emergency surgery, 12*, 5. <https://doi.org/10.1186/s13017-017-0117-6>
2. Costantini, T. W., Coimbra, R., Holcomb, J. B., Podbielski, J. M., Catalano, R., Blackburn, A., Scalea, T. M., Stein, D. M., Williams, L., Conflitti, J., Keeney, S., Suleiman, G., Zhou, T., Sperry, J., Skiada, D., Inaba, K., Williams, B. H., Minei, J. P., Privette, A., Mackersie, R. C., ... AAST Pelvic Fracture Study Group (2016). Current management of hemorrhage from severe pelvic fractures: Results of an American Association for the Surgery of Trauma multi-institutional trial. *The journal of trauma and acute care surgery, 80*(5), 717–725. <https://doi.org/10.1097/TA.0000000000001034>
3. Halawi M. J. (2015). Pelvic ring injuries: Emergency assessment and management. *Journal of clinical orthopaedics and trauma, 6*(4), 252–258. <https://doi.org/10.1016/j.jcot.2015.08.002>
4. Hsu, S. D., Chen, C. J., Chou, Y. C., Wang, S. H., & Chan, D. C. (2017). Effect of Early Pelvic Binder Use in the Emergency Management of Suspected Pelvic Trauma: A Retrospective Cohort Study. *International journal of environmental research and public health, 14*(10), 1217. <https://doi.org/10.3390/ijerph14101217>

Share your feedback

Please complete our survey to help make Queensland Trauma Education better

The survey should take no more than 5 minutes to complete.

Scan the QR code or visit:

[Evaluation Form - Clinical Skills Development Service \(csds.qld.edu.au\)](https://csds.qld.edu.au)



Queensland Trauma Education

Pelvic trauma – Haemodynamically unstable pelvic trauma: Immersive scenario – Facilitator resource kit

Published by the State of Queensland (Clinical Skills Development Service), 2024

Visit csds.qld.edu.au/qte

Email CSDS-Admin@health.qld.gov.au

Phone [+61 7 3646 6500](tel:+61736466500)

